

RENAL PATHOLOGY: NEPHROTIC SYNDROME

(Part II; 2026)

Astrid Weins, MD, PhD

Chief, Renal Pathology

Department of Pathology

Mass General Brigham

Assistant Professor of Pathology

Harvard Medical School





Astrid Weins, MD, PhD

aweins@bwh.harvard.edu

Chief, Renal Pathology, Mass General Brigham, Boston, MA

Assistant Professor of Pathology, Harvard Medical School

Renal pathologist, attending on the BWH diagnostic renal pathology service since 2011

I direct an active translational research program studying proteinuric kidney diseases

My research focuses on understanding the pathobiology of acute nephrotic syndrome and podocyte injury by integrating morphologic observations with imaging and tissue interrogation techniques

Disclosures (last 12 months)

- Patent and Licensure Fees

“Methods for identifying and treating patients with antibody-mediated acquired primary or recurrent idiopathic nephrotic syndrome”

Euroimmun GmbH

- Speaker fees

Genentech

- Advisory Board

Vera Therapeutics


- Funding

NIDDK, DoD

Objectives

- Brief review of glomerular structure
- Use a case-based discussion to highlight histopathologic findings in diffuse podocyte injury
- Correlate histopathologic findings with known etiologies in Minimal Change Disease and Membranous Nephropathy


Glomerular Structure - Recap



Cells: Endothelium
Epithelium (visceral and parietal)
Mesangium


Space: Capillary lumen
Urinary space

Matrix: Glomerular Basement Membrane
Mesangial Matrix



The podocyte –

a terminally
differentiated,
postmitotic
epithelial cell
enveloping the
glomerular
capillaries



SEM


TEM



High power cross section of the glomerular filtration barrier

Elements:

1. Fenestrated and charged endothelium
2. Dense GBM
3. Podocyte foot processes with slit diaphragms



Nephrin
Podocin
Neph1-3
CD2AP

ACTN4
Synaptopodin
Formins
Rac1

Integrins/Focal Adhesions Components of the GBM

TRPC6, TRPC5

Perico et al, Nat Rev Nephrol. 2016 Nov;12(11):692-710

**Diseases with direct and diffuse podocyte injury
and/or loss:**

“PRIMARY” PODOCYTOPATHIES

- 1. Minimal Change Disease**
- 2. Primary (Idiopathic) Focal and Segmental
Glomerulosclerosis**
- 3. Collapsing Glomerulopathy**
- (4. Membranous Nephropathy)**

Case 1

This 41-year-old man presents to the ED with chief complaint of a sore throat. A diagnosis of a URI is made.

Two weeks later, he notices swelling in his legs. This quickly progresses to total body swelling.

He is referred to your clinic by his PCP with 4+ proteinuria on dipstick. An UA shows a protein-to-creatinine ratio of 15 g/gCr.

He denies NSAID use, use of any other medication or use of illicit drugs. There is no significant past medical history.

Physical examination:

Unremarkable, except for a body weight of 237 lbs (up from 220 lbs 3 months ago), 2+ edema of LE

U/A :

1+ blood, 3+ protein; the sediment shows 0-5 RBC/hpf, 0 WBC/hpf, no cellular casts, few hyaline casts


Laboratory testing:

Hgb 17.7, WBC 7.1K, Plt 373; Na 139, K 4.6, Cl 107; Creatinine 1.5, BUN 14, glucose 90; Ca 7.6, TSH 6.93





ANA negative, ASO negative, ESR 121


An SPEP and UPEP are negative for paraproteins

A kidney biopsy was performed.




PAS







EM



EM



PAS




H&E

Diagnosis

Diffuse podocytopathy with minimal glomerular
changes
(Minimal Change Disease)

Acute Tubular Injury

Podocyte injury leads to generally reversible structural changes



(adapted from d'Agati et al., 2003)



The consequence of these alterations is severe albuminuria

Foot process effacement and loss of filtration surface

Focal defects in podocyte coverage

Increased hydraulic conductivity and ineffective reflection

→ **albuminuria**



Based on: Rennke HG and Denker BM. Renal Pathophysiology. The Essentials. Wolters Kluwer 2020.

Evidence for Autoimmunity




- **Corticosteroids** are effective at inducing remission, especially in children, yet relapse and steroid dependence are common, and in some patients, additional immunosuppressive agents are successfully used.
- **B-cell targeted therapies** (Rituximab) are effective in both adults and children with relapsing or steroid-dependent NS, suggesting a **potential autoantibody mediated etiology**.
- **Memory B cells** are selectively increased in patients with SSNS
- Further evidence for **B-cell involvement**: MCD was cured by measles, a disease that leads to B cell depletion and immune amnesia

Proposed antigenic podocyte targets in MCD

Antigen	Localization	Method of discovery	References
Ubiquitin Carboxyl Terminal Hydrolase L1 (UCHL1)	Cytoplasmic and nuclear (not expressed in normal glomeruli)	Proteomic Analysis	<i>Jamin et al, J Autoimmun 2018;</i> <i>Chebotareva et al, Front Med 2023</i>
Annexin A2 (ANXA2)	Cytoplasmic and membrane bound, phospholipid-binding	2D electrophoresis and Mass Spec	<i>Ye et al, Ann Transl Med, 2021</i>
Crumbs cell polarity complex component 2 (CRB2)	Junctional transmembrane/slit diaphragm	Mouse model only	<i>Hada et al, JASN 2022</i>
CD40	Cell membrane/ligand	Protein array (rFSGS)	<i>Delville et al, Sci Transl Med 2014;</i> <i>Chebotareva et al, 2023</i>
Nephrin	Transmembrane/slit diaphragm	Immunostaining and ELISA/IP	<i>Watts, Keller et al, JASN 2022</i> <i>Hengel et al, NEJM, 2024; KI 2025</i>
Podocin Neph1/Kirrel1	Transmembrane/slit diaphragm	STED, ELISA	<i>Raglianti et al, KI 2024, JASN 2025</i>

Evidence for IgG in our MCD biopsies


- Over many years, we have observed a **delicate extracapillary, punctate and diffuse glomerular staining for IgG** in ~64% of our MCD biopsies.




Widefield
Synpo
overlaid
with IgG
SIM

Punctate IgG staining in MCD biopsies colocalizes with nephrin


IgG (routine
epifluorescence)



IgG/nephrin (confocal)



SIM





Anti-nephrin antibodies are in circulation

We obtained **serum during active disease** and tested for **anti-nephrin by indirect ELISA** on some of our in-house patients.

All IgG+ MCD patients (MCD+) were **positive for anti-nephrin** antibodies.

In contrast, all non-nephrotic or nephrotic control patients, including 3 IgG- MCD patients, were serologically negative.

In all patients in whom we were also able to obtain a **remission** sample, anti-nephrin antibody levels were significantly **reduced**.




High % of anti-nephrin positivity in adults and kids

Hengel et al, NEJM 2024


B

Anti-Nephrin Prevalence in Adults with Glomerular Diseases




C

Anti-Nephrin Prevalence in Children with Idiopathic Nephrotic Syndrome




A

IP/ELISA Hybrid Assay




C

Proteinuria in Adults During Follow-Up




D






Proteinuria in Children During Follow-Up



E



Proof of causality in immunization and passive transfer models



Patient follow up

Several months later...

Chief complaints:


- Persistent lower extremity edema, otherwise normal physical exam
- Hyperlipidemia

Blood:


- Creatinine: 0.8 mg/dl
- BUN: 15 mg/dl
- Glucose: 85 mg/dl
- Serum Albumin: 3.1 mg/dl

Urine:

- Urine protein 3.6 g/24h
 - No hematuria
-
- All other tests and serologies were still negative



PAS




H&E

Cytokeratin stain


IHC

20 μ m




C3

IF




IgM


IF



EM



EM




Diagnosis


Progressive diffuse podocytopathy with early focal
and segmental glomerulosclerosis
?Collapsing features

All 3 morphological patterns can be seen in the same patient, at different time points!


MCD



FSGS




Collapsing glomerulopathy



It is important to establish:

1. Disease-initiating mechanisms (incl anti-nephrin)
2. Disease-perpetuating/progressing factors (environmental, genetic, other)
3. Predisposing factors (genetic background)

My concept of diffuse podocytopathies



Common diagnostic patterns observed in patients with Nephrotic Syndrome always affect the podocyte

“Podocytopathies”:

Minimal Change Disease

Focal Segmental Glomerulosclerosis

Collapsing Glomerulopathy

Membranous Nephropathy (idiopathic MN, Lupus)

Renal Amyloidosis

Nodular Glomerulosclerosis (Diabetic Nephropathy)

Case 3

A 63 yo man presents with “ankle swelling” which improves with a diuretic.

Heart: normal systolic function, LVEF 60%.

Hypercholesterolemia – started on simvastatin

No arthralgias, no hematuria.

Serology negative for HIV and Hepatitis.

Labs: BUN: 35 mg/dL; Cr: 0.92 mg/dL Alb: 2.5 g/dL Chol: 347 mg/dL


Urinalysis: 4+ protein/ +blood

24h urine: 9.7g protein/24hr


Urine sediment: Hyaline casts, oval fat bodies

Renal U/S: normal, 11cm left kidney; 11.8cm right kidney


A renal biopsy is performed.



PAS




PAS




IF

IgG




IF

C₃




This immunofluorescence image displays a glomerulus with prominent, bright green staining. The staining is distributed throughout the glomerular tuft, highlighting the capillary walls and the surrounding mesangial area. The pattern of staining is consistent with the deposition of C3, a component of the complement system, which is often seen in various forms of glomerulonephritis. The background is dark, indicating minimal staining in the surrounding tissue.




IgG1

This panel shows a fluorescence microscopy image of a tissue section stained with IgG1. The staining is localized and appears as a dense, irregular green mass against a black background.




IgG2

This panel shows a fluorescence microscopy image of a tissue section stained with IgG2. The staining is localized and appears as a dense, irregular green mass against a black background.




IgG3

This panel shows a fluorescence microscopy image of a tissue section stained with IgG3. The staining is localized and appears as a dense, irregular green mass against a black background.



IgG4


This panel shows a fluorescence microscopy image of a tissue section stained with IgG4. The staining is localized and appears as a dense, irregular green mass against a black background.



IF

Co-localization; IgG4-PLA2r



50 μm



Diagnosis

Membranous Nephropathy Stage II, PLA2r-positive


Membranous Nephropathy – explained!



- Process of immune complex deposition is understood and accepted.
- Specific epitopes are identified for the majority of antigens/cases.
- Immunofluorescence and serological tests are available.

“Lifting” of a podocyte by a growing deposit in early membranous nephropathy

Interference with podocyte attachment to the GBM




Membranous Nephropathy - Idiopathic no more!

Major **discoveries** of endogenous and exogenous antigens in last 2 decades, associated with various disorders

Presence of specific autoantibodies in the serum is **diagnostic for MN**, and predicts treatment response, disease activity and outcome

Slow disease **progression** and long **persistence** of immune complexes in the glomerulus leave a **long window** for both serological and histological detection



TARGETS TO REMEMBER IN MEMBRANOUS NEPHROPATHY

- M type phospholipase A2 receptor (PLA₂r)
- Exostosin 1/2 (EXT) (SLE)
- Neural epidermal growth factor-like1 prot (NELL1)
- Thrombospondin Type-1 Domain-Containing 7A (THSD7A)
- Semaphorin 3B
- alpha-enolase

PLA₂R > EXT > NELL1 > THSD7A and Semaphorin 3B

Congenital Membranous Nephropathy

- neutral endopeptidase (NEP)


Question 1

1) The target protein of the newly discovered and validated autoantibodies in minimal change disease is a component of the

- a) Glomerular basement membrane
- b) Podocyte actin cytoskeleton
- c) Slit diaphragm
- d) Focal adhesion complex

Answer 1

c) Nephrin is an essential component of the podocyte slit diaphragm.



Question 2

2) The most common autoantibody target in membranous nephropathy is

- a) NELL1
- b) Phospholipase A2 receptor 1
- c) dsDNA
- d) nephrin

Answer 2

b) Autoantibodies against the Phospholipase A2 receptor 1 account for 70-80% of cases of Membranous Nephropathy.

PLA2r > EXT > NELL1 > THSD7A and Semaphorin 3B

References

- Perico, L., Conti, S., Benigni, A. & Remuzzi, G. Podocyte-actin dynamics in health and disease. *Nat Rev Nephrol* **12**, 692-710 (2016).
- D'Agati, V. Pathologic classification of focal segmental glomerulosclerosis. *Semin Nephrol* **23**, 117-134 (2003).
- Watts, A.J.B., Keller K.H., et al. Autoantibodies against nephrin elucidate a novel autoimmune phenomenon in proteinuric kidney disease. *J Am Soc Nephrol* **33**(1):238-252 (2022).
- Hengel FE, Dehde S, Lasse M, et al. Autoantibodies targeting nephrin in podocytopathies. *N Engl J Med*. Published online May 25, 2024.
- Hengel FE, Dehde S, Yilmaz E, et al. Autoantibodies targeting nephrin in podocytopathies. *Kidney Int* 105, 1099-1103 (2025).
- Hoxha E, Reinhard L, Stahl RAK. *Nat Rev Nephrol*. Apr 28 (2022).
- Ronco P, Plaisier E, Debiec H. *J Clin Med* **10**(4):607 (2021).
- Beck, L.H., Jr., et al. M-type phospholipase A2 receptor as target antigen in idiopathic membranous nephropathy. *N Engl J Med* **361**, 11-21 (2009).
- Sethi, S., Madden, B. Mapping antigens of membranous nephropathy: almost there. *Kidney Int* **103**, 269-472 (2023).