

COVID-19 Vaccination in Immunocompromised Patients

Ann E. Woolley, MD, MPH
Associate Clinical Director of Transplant Infectious Diseases
Division of Infectious Diseases
Brigham and Women's Hospital
Instructor of Medicine
Harvard Medical School

Ann E. Woolley, MD, MPH



- Princeton University – BSE
- Icahn School of Medicine at Mount Sinai – MD, MPH
- Medicine Residency at Boston Medical Center
- Infectious Diseases Fellowship at MGH/BWH
- Transplant Infectious Diseases Fellowship at BWH
- Associate Clinical Director of Solid Organ Transplant Infectious Diseases
 - Clinical focus: transplant and oncology infectious diseases and cardiac device infections
 - Research focus: Clinical investigator, clinical trials




CDC: COVID-19 and Immunocompromised Patients

- *“Many conditions and treatments can cause a person to be immunocompromised or have a weakened immune system”*
 - Solid organ transplantation
 - Recent hematopoietic stem cell transplant
 - Active or recent treatment for solid tumor or hematologic malignancies
 - Severe primary immunodeficiency
 - HIV with a low CD4+ cell count or not on treatment
 - Prolonged use of corticosteroids, or use of other immune weakening medications (alkylating agents, antimetabolites, tumor-necrosis (TNF) blockers, and other biologic agents that are immunosuppressive or immunomodulatory)
- Having a weakened immune system may increase the risk of severe illness from COVID-19

Immune Response to SARS-CoV-2

Immune Responses Leading to Recovery or Death



Adequate immune responses

- Timely innate/adaptive responses
- Quick type 1 IFN response
- Activation of efficient antiviral response (clearance by macrophages)
- Activation of Th1 cells and B-cells for production of neutralizing antibodies

Inadequate immune responses

- Delayed/limited type 1 IFN
- Endothelial cell death
- Epithelial/endothelial leakage
- Overactivation/exhaustion T-cells and NK cells
- Accumulation of activated macrophages → cytokine storm


Immune-Related Risk Factors for Susceptibility to Severe COVID

- Older persons have impaired B-cell and T-cell immune responses vs younger persons
 - Impaired B-cell response: reduced ability to mount a neutralizing antibody response
 - Impaired T-cell response: we are learning more about the role of T-cell response in control of ongoing SARS-CoV-2
- Younger persons with impaired or dysregulated B-cell and T-cell immune responses may also be at risk

SARS-CoV-2 Impact on Immunocompromised Individuals

- Immunocompromised individuals comprise ~2.7% (~7 million) of U.S. adults
- More likely to get severely ill from COVID-19
- Higher risk for:
 - Prolonged SARS-CoV-2 infection and viral shedding
 - Viral evolution during infection and treatment (hospitalized patients)
- More likely to transmit SARS-CoV-2 to household contacts
- Lower antibody/neutralization titers to SARS-CoV-2 variants compared to immunocompetent individuals
- More likely to have vaccine breakthrough infections


Vaccination and Immunity



Vaccine Breakthrough Infections

- Lower vaccine effectiveness among immunocompromised individuals
 - 59-72% vaccine efficacy among immunocompromised vs. 90-94% among immunocompetent after 2nd dose of mRNA vaccine
- Immunocompromised individuals – *solid organ transplant recipients* – mount a less robust humoral response post-COVID vaccination compared to immunocompetent individuals


Percent of subjects with antibody response after two mRNA COVID-19 vaccine doses by immunocompromising condition and study (n=63)



- Studies that compared response after 1st and 2nd dose demonstrated less robust response after dose 1
- Antibody measurement and threshold levels vary by study protocol

Factors that most contributed to having a positive antibody?


- **Abdominal organ transplant** (kidney or liver) rather than a thoracic organ transplant (lung or heart)
- **>10 years out from transplant**
- ***Not* being on mycophenolate**



Vaccine Breakthrough Infections

- Lower vaccine effectiveness among immunocompromised individuals
 - 59-72% vaccine efficacy among immunocompromised vs. 90-94% among immunocompetent after 2nd dose of mRNA vaccine
 - Immunocompromised individuals – *solid organ transplant recipients* – mount a less robust humoral response post-COVID vaccination compared to immunocompetent individuals
 - Prior studies demonstrated an increased immune response post-3rd COVID vaccine dose when given 1-month post-2nd dose in transplant recipients
-

Randomized Trial of a 3rd Dose of Moderna Vaccine in Transplant Recipients (n=120)



RBD antibody (≥ 100 U/ml) 1
month post dose 3:

33 of 60 patients
(55%) vaccine group

vs.

10 of 57 patients
(18%) placebo group

COVID-19 Vaccination in Transplant Patients

- Many questions remain
 - What is the optimal timing of additional vaccine doses in immunocompromised individuals?
 - Does a mix-and-match strategy for mRNA COVID vaccines impact response?
 - What is the correlate for immune protection and does measuring SARS-CoV-2 spike Ab impact clinical care?
 - Objective of the BWH transplant COVID-19 vaccine study:
 - *Does a 3rd mRNA vaccine dose administered 6-months post-2nd dose enhance humoral immune response in transplant recipients?*
-

Single-center Prospective Cohort Study

Inclusion Criteria

- Heart, lung, kidney or dual organ txp recipient
- Had not received COVID vaccine prior to txp
- Did not have COVID-19 prior to receipt of 3rd vaccine dose

mRNA-1273 [Moderna]
BNT162b2 [Pfizer-BioNTech]

236 SOT
patients

Jan 2021

6 months

Feb 2022

- COVID-19 Infection?
- Survival


Roche Elecsys anti-SARS-CoV2
spike enzyme immunoassay

Proportion with a Positive Ab Post-3rd Vaccine Dose



	Positive Antibody Post-3 rd Dose	Negative Antibody Post-3 rd Dose
Total 236	132 (56%)	104 (44%)

Anti-RBD Antibodies Before and After 3rd Vaccine Dose



- <10 years post-transplant
- ▲ ≥10 years post-transplant

Heart
n = 48

Lung
n = 72

Kidney
n = 51

Multivariate Analysis



Characteristic	Adjusted Odds Ratio (95% CI)	Adjusted p-value
Age, >60 years old (<i>ref</i> = <60 yo)	0.38 (0.20 to 0.71)	0.0023
Years out from transplant (<i>ref</i> = 1 year)		
2 years	5.19 (1.42 to 18.99)	0.0129
3-5 years	5.83 (1.62 to 20.93)	0.0069
6-10 years	4.59 (1.25 to 16.86)	0.0216
11-15 years	10.96 (2.59 to 46.36)	0.0011
>= 16 years	16.61 (3.78 to 72.93)	0.0002
Organ type (<i>ref</i> = kidney)		
Lung	0.17 (0.07 to 0.39)	<0.0001
Heart	0.25 (0.11 to 0.59)	0.0013
Mycophenolate	0.29 (0.15 to 0.56)	0.0002


Outcomes	Positive Antibody (<i>n</i> = 132)	Negative Serology (<i>n</i> = 104)
COVID+ post-3 rd dose	30 (23%)	22 (21%)
Months between post-3 rd dose and COVID+	3.7 (3.0 – 4.3)	4.2 (1.3 – 4.6)
Hospitalized	6 (20%)	8 (36%)
Mortality	4 (3%)	4 (4%)

- Despite extended interval of 6 months, the proportion of non-responders (44%) was similar to other studies when the 3rd dose was given 1-month post-2nd vaccine dose
- Significant factors to having a positive spike antibody post-3rd vaccine dose
 - Age less than 60 years old
 - Having an abdominal organ transplant (kidney) rather than a thoracic organ transplant (heart > lung)
 - Being >10 years out from transplant
 - Not being on mycophenolate
- A similar proportion of individuals had COVID-19 infection post-3rd vaccine dose, irrespective of positive vs negative antibody result
 - Median time to COVID-19 infection was 4 months post-3rd vaccine dose

What additional strategies are there for those immunocompromised individuals who do not mount a robust humoral response to vaccination?


Strategies to increase the immunogenicity of COVID-19 vaccines in kidney transplant recipients

- RCT to compare 3 strategies with a control single dose mRNA-1273 vaccination in kidney transplant recipients (KTRs):
 - double vaccine dose, heterologous vaccination, and temporary discontinuation of mycophenolate mofetil or mycophenolic acid
- Repeated vaccination increases SARS-CoV-2-specific antibodies in KTRs, without further enhancement by use of a higher dose, a heterologous vaccine, or 2 weeks discontinuation of mycophenolate mofetil or mycophenolic acid



SARS-CoV-2 Viral Life Cycle

Neutralizing
Antibodies



Monoclonal Antibodies Bind to Spike Protein




Image credit: <https://www.nih.gov/>

Monoclonal antibodies

Bamlanivimab + Etesevimab


Casirivimab + Imdevimab

Sotrovimab

Bebtelovimab

Tixagevimab/cilgavimab – COVID-19 Pre-exposure Prophylaxis

Figure 1 Kaplan Meier: Cumulative Incidence of Symptomatic COVID-19* (PROVENT)



Levin. IDWeek 2021. Abstr LB5.


(A)

All SOTRs

COVID-19 (%)

Log-rank
 $P < 0.001$

Pemivibart 2024



Question #1

I am an adult in 2024 who underwent a stem cell transplant 6 months ago and have had one or more doses of a COVID-19 vaccine in the past. What is the current recommendation?

- A. Get a first dose of the updated 2023-2024 COVID mRNA vaccine
 - B. Get a second dose 3 weeks after the last dose
 - C. Get a third dose at least 8 weeks after the last dose
 - D. Get one dose only of the updated 2023-2024 COVID vaccine
 - E. It depends
 - F. A, B, C
-

Question #1

I am an adult in 2024 who underwent a stem cell transplant 6 months ago and have had one or more doses of a COVID-19 vaccine in the past. What is the current recommendation?

- A. Get a first dose of the updated 2023-2024 COVID mRNA vaccine
 - B. Get a second dose 3 weeks after the last dose
 - C. Get a third dose at least 8 weeks after the last dose
 - D. Get one dose only of the updated 2023-2024 COVID vaccine
 - E. It depends
 - F. A, B, C
-

Question #2

Vaccinated immunocompromised individuals no longer have an increased risk of hospitalization and death from COVID-19 compared to non-immunocompromised individuals.

- A. True
 - B. False
-

Question #2

Vaccinated immunocompromised individuals no longer have an increased risk of hospitalization and death from COVID-19 compared to non-immunocompromised individuals.

A. True

B. False

TABLE 3. Association of vaccination status* with intensive care unit admission and in-hospital death among patients hospitalized for COVID-19, by immunocompromise status—COVID-NET, 10 states, † March 1, 2021–February 28, 2022

Vaccination status*	No. (weighted %) [§]											
	Immunocompromised†						Not immunocompromised**					
	ICU admission			Death			ICU admission			Death		
	Yes	No	aOR (95% CI)	Yes	No	aOR (95% CI)	Yes	No	aOR (95% CI)	Yes	No	aOR (95% CI)
Vaccinated	85 (25.0)	269 (75.0)	1.01 (0.64–1.58)	55 (16.5)	298 (83.5)	1.34 (0.71–2.51)	257 (18.7)	1,044 (81.3)	0.85 (0.60–1.12)	113 (9.5)	1,188 (90.5)	0.58 (0.39–0.86) ^{††}
Unvaccinated	129 (25.5)	351 (74.5)	Ref	66 (12.9)	413 (87.1)	Ref	1,121 (21.6)	3,771 (78.4)	Ref	488 (10.1)	4,409 (89.9)	Ref

Thank you!

Ann E. Woolley, MD, MPH
awoolley@bwh.harvard.edu



References

1. Al Jurdi A, Morena L, Cote M, Bethea E, Azzi J, Riella LV. Tixagevimab/cilgavimab pre-exposure prophylaxis is associated with lower breakthrough infection risk in vaccinated solid organ transplant recipients during the omicron wave. *Am J Transplant*. 2022 Jun 21. doi: 10.1111/ajt.17128. Epub ahead of print. PMID: 35727916.
2. Kim L, Garg S, O'Halloran A, et al. Risk factors for intensive care unit admission and in-hospital mortality among hospitalized adults identified through the US coronavirus disease 2019 (COVID-19)–associated hospitalization surveillance network (COVID-NET). *Clin Infect Dis* 2021;72:e206–14. <https://doi.org/10.1093/cid/ciaa1012> PMID:32674114
3. Garg S, Patel K, Pham H, et al. Clinical trends among U.S. adults hospitalized with COVID-19, March to December 2020: a cross-sectional study. *Ann Intern Med* 2021;174:1409–19. <https://doi.org/10.7326/M21-1991> PMID:34370517
4. Tenforde MW, Patel MM, Gaglani M, et al.; IVY Network. Effectiveness of a third dose of Pfizer-BioNTech and Moderna vaccines in preventing COVID-19 hospitalization among immunocompetent and immunocompromised adults—United States, August–December 2021. *MMWR Morb Mortal Wkly Rep* 2022;71:118–24. <https://doi.org/10.15585/mmwr.mm7104a2> PMID:35085218
5. Tenforde MW, Self WH, Gaglani M, et al.; IVY Network. Effectiveness of mRNA vaccination in preventing COVID-19–associated invasive mechanical ventilation and death—United States, March 2021–January 2022. *MMWR Morb Mortal Wkly Rep* 2022;71:459–65. <https://doi.org/10.15585/mmwr.mm7112e1> PMID:35324878
6. Harpaz R, Dahl RM, Dooling KL. Prevalence of immunosuppression among US adults, 2013. *JAMA* 2016;316:2547–8. <https://doi.org/10.1001/jama.2016.16477> PMID:27792809
7. Chow N, Fleming-Dutra K, Gierke R, et al.; CDC COVID-19 Response Team. Preliminary estimates of the prevalence of selected underlying health conditions among patients with coronavirus disease 2019—United States, February 12–March 28, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:382–6. <https://doi.org/10.15585/mmwr.mm6913e2> PMID:32240123
8. Andrejko KL, Pry JM, Myers JF, et al.; California COVID-19 Case-Control Study Team. Effectiveness of face mask or respirator use in indoor public settings for prevention of SARS-CoV-2 infection—California, February–December 2021. *MMWR Morb Mortal Wkly Rep* 2022;71:212–6. <https://doi.org/10.15585/mmwr.mm7106e1> PMID:35143470
9. Sami S, Horter L, Valencia D, et al. Investigation of SARS-CoV-2 transmission associated with a large indoor convention—New York City, November–December 2021. *MMWR Morb Mortal Wkly Rep* 2022;71:243–8. <https://doi.org/10.15585/mmwr.mm7107a4> PMID:35176005
10. Wiltz JL, Feehan AK, Molinari NM, et al. Racial and ethnic disparities in receipt of medications for treatment of COVID-19—United States, March 2020–August 2021. *MMWR Morb Mortal Wkly Rep* 2022;71:96–102. <https://doi.org/10.15585/mmwr.mm7103e1> PMID:35051133
11. Fu J, Reid SA, French B, et al.; COVID-19 and Cancer Consortium (CCC19). Racial disparities in COVID-19 outcomes among black and white patients with cancer. *JAMA Netw Open* 2022;5:e224304. <https://doi.org/10.1001/jamanetworkopen.2022.4304> PMID:35344045
12. Singson JR, Kirley PD, Pham H, et al. Factors Associated with Severe Outcomes Among Immunocompromised Adults Hospitalized for COVID-19 — COVID-NET, 10 States, March 2020–February 2022. *MMWR Morb Mortal Wkly Rep* 2022;71:878–884. DOI: <http://dx.doi.org/10.15585/mmwr.mm7127a3>

