



Brigham and Women's Hospital
Founding Member, Mass General Brigham

ACUTE KIDNEY INJURY SYNDROMES

Alice M. Sheridan, M.D.
Associate Physician
Division of Nephrology
Brigham and Women's Hospital
Assistant Professor of Medicine
Harvard Medical School



Alice M. Sheridan, M.D.



New York Medical College
Medicine Residency, Boston Medical Center
Renal Fellowship, Boston Medical Center
Assistant Professor of Medicine, Harvard Medical
School

DISCLOSURES

I have no relevant financial relationships with ineligible companies.



OBJECTIVES

- To define the common syndromes that underlie acute kidney injury (AKI) and recognize limitations of definitions
- To describe the clinical presentation and prognosis of select syndromes



AKI definition

- A group of syndromes characterized by an acute decrease in glomerular filtration.
- The degree of reduction in glomerular filtration and/or urine output provide the sole criteria for diagnosis.
- Acute Dialysis Quality Initiative (2004) and revised by the AKI network (2007) and KDIGO (2012)
- KDIGO new guidelines? (Draft issued in March 2026)



KDIGO

- An increase in serum creatinine by > 0.3 mg/dL (>26.5 micromol/L) within 48 hours, or
- An increase in serum creatinine to > 1.5 times baseline that is known or presumed to have occurred within the past 7 days, or
- Urine volume <0.5 mL/kg/hour for six hours.
- Increase in serum cystatin C by >1.5 times baseline that is known or presumed to have occurred within the past 7 days. KDIGO DRAFT 2026.
- AKD – AKI by creatinine/cystatin criteria or decline in eGFR <60 mL/min per 1.73m^2 or decrease in eGFR by 35mL/min from baseline or an increase in serum creatinine $>50\%$ or structural AKI or emergence of albuminuria, hematuria or leukocyturia for duration <3 months.
- Practice point – incorporates structural abnormalities (ie biomarkers, (TIMP-2, IGFBP7, NGAL, KIM-1, LFBP. (Draft)



Urine output

- Not easily measured outside of the ICU setting
- Imparts prognostic information
- Intraoperative oliguria predicts AKI in high risk patients
- Early identification of AKI allows effective intervention (PrevAKI study)
- Real time monitoring (Accuryn Monitoring System) allows early identification in cardiac surgery patients



Cystatin C more accurate in AKI

- Use cystatin C when creatinine expected to be less accurate
 - muscle wasting, neurologic illness, spinal cord, AKA, interfering drugs
- Better correlation with iohexol over length of ICU stay (Haines CJASN 18:997-1005, 2023)
- Earlier detection of AKI (12-24 hours compared with 12 to 72 hours for creatinine) (Hooper CJASN 20: 477-484, 2025)
 - Smaller volume of distribution (extracellular versus TBW)
 - Faster time to doubling
- Best predictor of AKI (Lan Medicine 102:40. 2023)
- Decreases more quickly in recovering patients (Gharaibeh Kidney Int Rep 3: 337-342, 2018)



Epidemiology and outcomes

- High resource settings, AKI occurs in approximately 20% adult hospitalized patients
- AKI is risk factor for mortality among patients with sepsis;
- 3-5X increase in mortality at 60 days.
- Early reversal of AKI associated with 1 year survival is >90%
- No reversal associated with <40% survival.



Community Acquired (CA-AKI) versus Hospital-Acquired (HA-AKI)

- Different risk factors, epidemiology and outcome
- CA-AKI- present at time of admission or recognized shortly thereafter. Results from factors present in the community
- HA-AKI – Develops after 48 hours. Due to therapeutic or diagnostic intervention or a sequela of another illness
- Most studies and clinical practice guidelines pertain to HA-AKI



What are the common underlying syndromes

- Prerenal/acute tubular necrosis
- Hepatorenal
- Cardiorenal
- Rapidly progressive glomerulonephritis
- Acute interstitial nephritis
- Crystalopathies
- Obstruction



AKI-EPI Study- HA-AKI

- Prerenal/acute tubular necrosis
 - Sepsis 41%
 - Hypovolemia 34%
- Hepatorenal 3.2%
- Cardiorenal
 - Cardiogenic shock 13%
- Rapidly progressive glomerulonephritis
- Acute interstitial nephritis
 - “Drug related” – 14%
- Crystalopathies
- Obstruction (1.4%)

Hoste et al, Intensive Care Medicine 2015 41: 1411



What about CA-AKI?

- Most cases of CA-AKI are in developing countries of low- or middle income
 - Infection
 - Volume depletion from diarrhea illness
 - Poisoning, animal bites, venomous stings
 - Poor obstetrical care
 - Heat illness



2016 Meta-analysis

- Incidence in low or middle income countries similar to high income countries.
- Increased mortality and CKD/ESKD risk
- High income – risk factors diabetes, CVD, CKD, older age
- Lower/middle income – younger age, few comorbidities



What about CA-AKI in high income or developed countries?

- Not well studied
- Outpatient population
- Incidentally detected
- Duration is not known
- Role of biopsy



Case 1

- 65-year-old man CKD eGFR 40-55 mL/min per 1.73m² status post bilateral lung transplant and likely chronic calcineurin toxicity.
- Presented with increased creatinine that occurred over the course of 3 months. Tacrolimus level was therapeutic throughout.
- He had diarrhea attributed to small intestine bacterial overgrowth (SIBO) , with significant weight loss (30 lbs) . Treated with Augmentin.
- Ultrasound showed nonobstructing right nephrolithiasis, right renal cyst. Kidney size 12.1, 12.3 cm




Case 1

- At the time of your visit, the diarrhea has abated and he has stopped antibiotics.
- His diet is back to normal except that he is now drinking spinach smoothies and taking vitamin C gummies (1-2 daily, but occasionally takes as many as 3-4) in effort to recover his previous health status.
- He is euvolemic on exam. His creatinine remains elevated at 3.54 mg/dL.



Case 1

- You elect to do a kidney biopsy.
- Light microscopy showed widespread tubular injury with distended lumina and many intraluminal birefringent crystals under polarized light.
- There was interstitial fibrosis but no active cellular infiltrate. A representative micrograph is shown here.



(Micrograph and biopsy interpretation provided by Dr. Helmut R. Rennke, Renal Pathology, Brigham and Women's Hospital. Boston MA)

What is the likely cause of his persistent AKI?

- A. Prolonged ischemic acute tubular injury from volume depletion
- B. Drug induced acute interstitial nephritis
- C. Oxalate nephropathy
- D. Acute phosphate nephropathy



What is the likely cause of his persistent AKI?

- A. Prolonged ischemic acute tubular injury from volume depletion
- B. Drug induced acute interstitial nephritis
- C. Oxalate nephropathy
- D. Acute phosphate nephropathy



Case 1 continued

A. Prolonged ischemic acute tubular injury from volume depletion

The tubular injury observed on biopsy is commonly observed with oxalate deposition.

B. Drug induced acute interstitial nephritis Although interstitial nephritis may be observed with oxalate nephropathy, this patient did not have an active interstitial cellular infiltrate as would be expected with drug induced interstitial nephritis.

D. Acute phosphate nephropathy Calcium phosphate crystals are not birefringent under polarizing light, and unlike calcium oxalate, are identified with von Kossa stain. (also no risk factors in the history)



Oxalate nephropathy

- Secondary hyperoxaluria results from prolonged antibiotics and excessive intake of dietary oxalate and the oxalate precursor, ascorbic acid, in the setting of volume depletion.
- Oral antibiotics alter the bowel flora with decreases in both *Oxalobacter formigenes* (which degrades dietary oxalate) and butyrate-forming bacteria (which maintain the intestinal mucosa and regulate intestinal oxalate transporters)
- Increased systemic oxalate absorption and hyperoxaluria cause nephrolithiasis and oxalate nephropathy . Widespread oxalate deposition strongly supports oxalate nephropathy.



Secondary Hyperoxaluria and nephrolithiasis

- Major risk factor for calcium oxalate stones
- Meta-analysis of 12 studies increased risk of stones after Roux-en-Y (RR 1.79, 95% CI, 1.54-2.10)
- Multiple studies demonstrate increased stone risk after bariatric surgery
- Multiple studies demonstrate increased stone risk with Crohn's , bowel resections, IBD, ulcerative colitis



Secondary Hyperoxaluria –nephropathy (oxalosis)

- Less well documented – Requires biopsy. Easier to count kidney stones
- Systematic review and metanalysis (n-108)
 - 88% fat malabsorption
 - 20% dietary oxalate
 - Over half required dialysis
 - No complete recovery
 - 42 % partial recovery
 - Excluded patients with short duration of exposure (ethylene glycol star fruit)
 - Most common presentation AKI (35%) or CKD+AKI (29%)



Secondary Hyperoxaluria –Treatment

- Fluids (2.5 liters urine daily)
- Low oxalate diet
- Low fat diet
- Increased oral calcium
- Citrate






Case 2

- This patient presented to his primary care physician with nausea and vomiting and found to have creatinine of 6.1 mg/dL. Urinalysis showed 4+ heme
- Four days prior he had developed calf pain during hike). Urine turned dark after onset of pain. He took 400 mg ibuprofen during the hike
- He has had multiple episodes exactly like this (10-12 over past ten years).



Case 2-Biopsy



- Tubular injury with luminal dilatation, loss of brush border and flattened epithelial cells .
- Tubules contained myoglobin casts.
- Interstitium contained an inflammatory infiltrate of lymphocytes and plasma cells, neutrophils and eosinophils .

What was the most likely cause of ATN?

- A) Rhabdomyolysis from vigorous exercise
- B) Neuromuscular disorder
- C) Hypokalemia
- D) NSAIDs



What was the most likely cause of ATN?

- A) Rhabdomyolysis from vigorous exercise
- B) Neuromuscular disorder
- C) Hypokalemia
- D) NSAIDs



Rhabdomyolysis

- Crush syndrome, limb compression
- Strenuous exercise, seizures
- Disorders of glycolysis, glycogenolysis, lipid metabolism
- Infections (Influenza A and B, COVID), coxsackie-virus, HIV, EBV, Strep, Staph,
- Heat stroke, malignant hyperthermia, malignant neuroleptic syndrome
- Hypokalemia, hypophosphatemia, hypocalcemia, DKA
- Drugs: fibrates, statins, alcohol, heroin, cocaine



Neuromuscular causes of rhabdomyolysis

- McArdle's disease
- Carnitine palmitoyl transferase deficiency
- Muscular dystrophies (Becker, limb girdle)
- Clinically evident ATN is uncommon
- Chronic tubulointerstitial nephritis has been described in patients with McArdle's , and other myopathies



Case 3

- 50 year old man referred for increased creatinine (1.1–1.48 mg/dL) noted over 3 months in setting of starting ACE inhibitor. ACE inhibitor stopped. Repeat creatinine stable at 1.40 to 1.50 mg/dL
- PMH remarkable for new onset hypertension and recent diagnosis of osteoporosis with vertebral fracture
- Occasional NSAID use, No PPI use



Case 3

- Urinalysis showed trace wbc on one occasion, sediment no casts, no crystals
- microalbumin/creatinine ratio normal
- SPEP, light chains negative, total /ionized calcium normal
- ANCA, ANA, dsDNA, anti-Ro/anti-La negative
C3,C4 normal




Presentation is most consistent with...

- A. Drug-induced interstitial nephritis
- B. Sarcoidosis
- C. TINU syndrome
- D. Infection-related interstitial nephritis



Biopsy



Chronic interstitial nephritis with a single perivascular granuloma

(Micrograph and biopsy interpretation provided by Dr. Helmut R. Rennke, Renal Pathology, Brigham and Women's Hospital. Boston MA)

Acute Interstitial Nephritis

- Common cause of unexplained AKI
- Drug related in >70 %
- Infections (legionella, leptospirosis, CMV, streptococcus)
- SLE, Sjogren's disease, sarcoidosis
- TINU (tubulointerstitial nephritis + uveitis) syndrome



Granulomatous Interstitial Nephritis

- Less common form of interstitial nephritis
- Drug related
- Infection - Mycobacterium, fungus (histoplasmosis, coccidiomycosis), bacteria, spirochetes, parasites.
- Tubulointerstitial nephritis and uveitis (TINU)
- Sarcoidosis
- Granulomatosis and polyangiitis (GPA)



Evaluation

- History
- ANCA, ACE, ANA, C3, C4, dsDNA
- Look for Infection - Mycobacterium, fungus (histoplasmosis, coccidiomycosis), bacteria, spirochetes, parasites.
- Tubulointerstitial nephritis and uveitis (TINU)
- Sarcoid – Imaging (CXR, CT, bronchoscopy)
 - 23% sarcoidosis patient have history of major osteoporosis fracture
 - Increased risk of vertebral fractures even off glucocorticoids



Case 4

- An 81 year old woman presents with anuria, nausea and vomiting 3 days after receiving zoledronate infusion.
- Creatinine 10 mg/dL (up from 1.2 mg/dL).
- Her creatinine declines on the third day and she does not require RRT.



Case 4

- You are asked by the patient and her endocrinologist about the risks of treating osteoporosis in patients with underlying CKD.
-
- Which of the following statements are true about bisphosphonates, and zoledronate in particular:



Case-4

- A) The risk of AKI is high with zoledronate and palmidronate even in the absence of CKD
- B) Underlying CKD did not increase her risk of AKI
- C) AKI is an unlikely but well described side effect of zoledronate. While not contraindicated, it is most prudent to avoid further use in this patient.
- D) The most common histologic lesion associated with zoledronate is collapsing glomerulopathy



Case 4

- A) The risk of AKI is high with zoledronate and palmidronate even in the absence of CKD
- B) Underlying CKD did not increase her risk of AKI
- C) AKI is an unlikely but well described side effect of zoledronate. While not contraindicated, it is most prudent to avoid further use in this patient.
- D) The most common histologic lesion associated with zoledronate is collapsing glomerulopathy



Bisphosphonates and AKI

- Pamidronate (high dose) - collapsing focal and segmental glomerulosclerosis (FSGS) in 2001 .
- Zoledronic acid (monthly infusions for myeloma or Paget's disease - AKI in 2003
- Nephrotoxicity of IV bisphosphonates is dose dependent and infusion rate dependent, and may be reduced by extending the time interval between infusions.



TAKE HOME MESSAGES

- Criteria that define AKI are imperfect
- Prognostic importance of urine output increasingly appreciated.
- Cystatin C is a better marker than creatinine
- Multiple conditions underlie AKI. Biopsy is commonly required.



REFERENCES-AKI

Hoste EA, et al. Epidemiology of acute kidney injury in critically ill patients: the multinational AKI-EPI study. *Intensive Care Med.* 2015;41(8):1411-23.

James MT, et al. Incidence and Prognosis of Acute Kidney Diseases and Disorders Using an Integrated Approach to Laboratory Measurements in a Universal Health Care System. *JAMA Netw Open.* Apr 5 2019;2(4):e191795. 2019.1795

Priyanka P, et al. The impact of acute kidney injury by serum creatinine or urine output criteria on major adverse kidney events in cardiac surgery patients. *The Journal of thoracic and cardiovascular surgery.* 2021;162(1):143-151 e147.

Bianchi NA, et al. Association of Oliguria With Acute Kidney Injury Diagnosis, Severity Assessment, and Mortality Among Patients With Critical Illness. *JAMA Netw Open.* 2021;4(11):e2133094



REFERENCES-AKI

- Pang Z, et al. The correlation of intraoperative oliguria with acute kidney injury after noncardiac surgery: a systematic review and meta-analysis. *Int J Surg*. 2023; 109(3):449
- Milder, D. Association between intraoperative oliguria and postoperative acute kidney injury in non-cardiac surgical patients: a systematic review and meta-analysis. *J Anesth* 2023 Apr;37(2):219-233.
- Ronco C, et al. Acute kidney injury. *The Lancet* 2019; 394: 1949
- Hoste EAJ, et al. Global epidemiology and outcomes of acute kidney injury. *Nat Rev Nephrol*. 2018; 14: 607.
- Mehta RL, et al. International Society of Nephrology's Oby25 initiative for acute kidney injury (zero preventable deaths by 2025): a human rights case for nephrology. *Lancet*. 2015 Jun 27;385(9987):2616-43. doi: 10.1016/S0140-6736(15)60126-X. Epub 2015 Mar
- Hooper L et al. The Kinetics of Cystatin C and Serum Creatinine in AKI. *Clin J Am Soc Nephrol*. 2025; 20: 477-484



REFERENCES-ATN

McCarron DA, et al . Chronic tubulointerstitial nephritis caused by recurrent myoglobinuria. Arch Int Med 1980;140:1106-7

James MT, Levey AS, Tonelli M, et al. Incidence and Prognosis of Acute Kidney Diseases and Disorders Using an Integrated Approach to Laboratory Measurements in a Universal Health Care System. JAMA Netw Open. 2019;2(4):e191795.



REFERENCES-Hyperoxaluria

McCarron DA, et al . Chronic tubulointerstitial nephritis caused by recurrent myoglobinuria. Arch Int Med 1980;140:1106-7

James MT, Levey AS, Tonelli M, et al. Incidence and Prognosis of Acute Kidney Diseases and Disorders Using an Integrated Approach to Laboratory Measurements in a Universal Health Care System. JAMA Netw Open. 2019;2(4):e191795.



REFERENCES-AIN

McCarron DA, et al . Chronic tubulointerstitial nephritis caused by recurrent myoglobinuria. Arch Int Med 1980;140:1106-7

James MT, Levey AS, Tonelli M, et al. Incidence and prognosis of acute kidney diseases and disorders using an integrated approach to laboratory measurements in a universal health care system. JAMA Netw Open. 2019;2(4):e191795.

Caffarelli C. et al. Osteoporosis and major fragility fractures (MOF) in sarcoidosis patients: association with disease severity. Aging Clinical and Experimental Research 2023 35:3015-3022

Bours S. Risk of vertebral and non-vertebral fractures in patients with sarcoidosis: a population-based cohort. Osteoporosis Int 2016; 27:1630-1610

